



Alabama

Live births, abortions, and pregnancies

Live birth rates, 1982–1992



Live birth rates,* 1991



Pregnancy rates, 1990

Pregnancy rates are not available because Alabama did not report abortion data to CDC.

National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
8th highest	15th highest	12th highest

*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




Sexually transmitted diseases Alabama

Gonorrhea rates, 1982–1992



Gonorrhea rates, 1992



National gonorrhea rate ranking,* 1992

Males – 3rd highest Females – 5th highest



*National ranking refers to males and females aged 10-24 years.



Chlamydia cases and rates, 1988–1992

Chlamydia data are not available for Alabama.

Acquired immunodeficiency syndrome (AIDS) cumulative cases,* 1981–1992




*Missing bar indicates fewer than six cumulative cases.



Related risk behaviors,* 1993


Alabama

Had sexual intercourse (lifetime)




Question not included in survey.

Had sexual intercourse with four or more people (lifetime)




Question not included in survey.

Had sexual intercourse during the past three months (currently sexually active)




Question not included in survey.

Used a condom during last sexual intercourse (among currently sexually active)




Question not included in survey.

Used birth control pills during last sexual intercourse (among currently sexually active)



Question not included in survey.

Injected illegal drugs (lifetime)



*Related risk behaviors refer to high school students.
Sources: CDC—NCHS (birth data), NCPS (gonorrhea data), NCID (AIDS data); 1993 Alabama Youth Risk Behavior Survey (risk behavior data).

